

Northwest 6320 N. La Cholla Blvd., #200 Tucson, AZ 85741 (520) 382-8200 Oro Valley 12315 N. Vistoso Park Rd. Oro Valley, AZ 85755 (520) 544-9700 St. Mary's 395 N. Silverbell Rd., #101 Tucson, AZ 85745 (520) 382-8200

POSTOPERATIVE INSTRUCTIONS: ELBOW SURGERY

FROM HOSPITAL TO HOME

- You must be accompanied by an adult who can drive you home after surgery.
- * You may have some dizziness from the effects of anesthesia and pain medications.
- * You cannot drive, operate machinery, drink alcohol or make critical decisions for at least 24 hours.

SURGICAL DRESSING

- ❖ You will be in a splint for your elbow until your first postoperative visit.
- ❖ You must keep the splint clean and dry.
- ❖ If you would like to shower, you need to ensure that your splint remains dry.
- ❖ At your first postoperative visit your sutures will be removed.

MEDICATIONS

- A narcotic pain medication will be prescribed to take as needed- please read and follow the instructions on the bottle.
- ❖ If you had a regional block performed by anesthesia, consider taking pain medication prior to the block wearing off completely for better control.
- Nausea and/or vomiting is a side effect of narcotic medication- if this should happen, consider changing to liquid diet and decreasing activity level.
- Constipation is another side effect of narcotic medication- you will also have a prescription for a stool softener.
- ❖ While taking narcotic pain medication you cannot drink alcohol or drive.
- Try to wean off the narcotic pain medication as soon as your pain allows. You can transition to over-the-counter Tylenol (acetaminophen) as needed.

ACTIVITY

❖ Your arm will be in a sling and should remain in the sling until your first postoperative visit- take out for hygiene and to periodically move your wrist and fingers.

APPOINTMENTS

- ❖ You will see Dr. Kim 2 weeks after surgery.
- For questions or concerns, please contact the surgery scheduler directly or call 520-784-6200.
- ❖ You should schedule a physical therapy appointment to begin approximately 2-4 weeks after surgery. The timing will be discussed during your postoperative visit.